

# Effect of Vitamin D<sub>3</sub> Supplementation vs Placebo on Hospital Length of Stay in Patients with Severe COVID-19: A Multicenter, Double-blind, Randomized Controlled Trial

**Short title:** Vitamin D<sub>3</sub> Supplementation in COVID-19

Igor H. Murai, PhD<sup>\*</sup>; Alan L. Fernandes, PhD<sup>\*</sup>; Lucas P. Sales, MSc; Ana J. Pinto, BSc; Karla F. Goessler, PhD; Camila S. C. Duran, MD; Carla B. R. Silva, MD; André S. Franco, MD; Marina B. Macedo, MD, MSc; Henrique H. H. Dalmolin, MD; Janaina Baggio, MD; Guilherme G. M. Balbi, MD; Bruna Z. Reis, PhD; Leila Antonangelo, MD, PhD; Valeria F. Caparbo, PhD; Bruno Gualano, PhD; Rosa M. R. Pereira, MD, PhD.

<sup>\*</sup>These authors contributed equally

**Manuscript word count:** 2,605 words.

**Corresponding author (to whom reprint requests should be addressed):**

Rosa Maria Rodrigues Pereira, MD, PhD.  
Rheumatology Division, Faculdade de Medicina FMUSP, 3º andar, Universidade de Sao Paulo, Sao Paulo, SP, BR.  
Av. Dr. Arnaldo, 455 – Pacaembu - São Paulo, SP – Brasil  
Postal code: 01246-903  
Phone: + 55 11 3061.7490 e-mail: rosamariarp@yahoo.com

**NOTE:** This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

**Key points:**

**Question:** Can vitamin D<sub>3</sub> supplementation reduce hospital length of stay in hospitalized patients with severe COVID-19?

**Findings:** In this double-blind, randomized, placebo-controlled trial involving 240 hospitalized patients with severe COVID-19, a single dose of 200,000 IU of vitamin D<sub>3</sub> supplementation was safe and effective in increasing 25-hydroxyvitamin D levels, but did not significantly reduce hospital length of stay (hazard ratio, 1.12) or any other clinically-relevant outcomes compared with placebo.

**Meaning:** Vitamin D<sub>3</sub> supplementation does not confer therapeutic benefits among hospitalized patients with severe COVID-19.

**Importance:** Patients with COVID-19 may exhibit 25-hydroxyvitamin D deficiency, but the beneficial effects of vitamin D<sub>3</sub> supplementation in this disease remain to be proven by randomized controlled trials.

**Objective:** To investigate the efficacy and safety of vitamin D<sub>3</sub> supplementation in patients with severe COVID-19.

**Design, Setting, and Participants:** This is a multicenter, double-blind, randomized, placebo-controlled trial conducted in two centers (a quaternary hospital and a field hospital) in Sao Paulo, Brazil. The trial included 240 hospitalized patients with severe COVID-19. The study was conducted from June 2, 2020 to October 7, 2020.

**Interventions:** Patients were randomly allocated (1:1 ratio) to receive either a single oral dose of 200,000 IU of vitamin D<sub>3</sub> or placebo.

**Main Outcomes and Measures:** The primary outcome was hospital length of stay, defined as hospital discharge from the date of randomization or death. Secondary outcomes were mortality, admission to ICU, mechanical ventilation requirement, and serum levels of 25-hydroxyvitamin D, creatinine, calcium, C-reactive protein, and D-dimer.

**Results:** Of 240 randomized patients (mean age, 56 years; 56% men), 232 (96.7%) were included in the primary analysis. Log-rank test showed that hospital length of stay was comparable between the vitamin D<sub>3</sub> supplementation and placebo groups (7.0 days [95% CI, 6.1 to 7.9] and 7.0 days [95% CI, 6.2 to 7.8 days]; hazard ratio, 1.12 [95% CI, 0.9 to 1.5];  $P = .379$ ; respectively). The rate of mortality (7.0% vs 5.1%;  $P = .590$ ), admission to ICU (15.8% vs 21.2%;  $P = .314$ ), and mechanical ventilation requirement (7.0% vs 14.4%;  $P = .090$ ) did not significantly differ between groups. Vitamin D<sub>3</sub> supplementation significantly increased serum 25-hydroxyvitamin D levels compared to

1 placebo (difference, 24.0 ng/mL [95% CI, 21.0% to 26.9%];  $P = .001$ ). No adverse  
2 events were observed.

3 **Conclusions and Relevance:** Among hospitalized patients with severe COVID-19,  
4 vitamin D<sub>3</sub> supplementation was safe and increased 25-hydroxyvitamin D levels, but  
5 did not reduce hospital length of stay or any other relevant outcomes vs placebo. This  
6 trial does not support the use of vitamin D<sub>3</sub> supplementation as an adjuvant treatment of  
7 patients with COVID-19.

8 **Trial Registration:** ClinicalTrials.gov Identifier: NCT04449718

9

# Introduction

A growing body of evidence has indicated that vitamin D may enhance the innate<sup>1-3</sup> and adaptive immunity.<sup>4, 5</sup> Since antigen-presenting cells have the ability to synthesize 1,25-dihydroxyvitamin D (the active form of vitamin D) from 25-hydroxyvitamin D, it has been postulated that vitamin D supplementation could improve the function of macrophages and dendritic cells, thereby ameliorating overall immune response.<sup>6</sup> In fact, insufficient vitamin D status has been suggested as a potential risk factor for non-communicable<sup>7</sup> and acute respiratory tract diseases,<sup>8, 9</sup> including viral infections.<sup>10</sup> In this context, it has been recently conjectured that optimal levels of vitamin D could play important immunomodulatory and anti-inflammatory roles, thereby benefiting patients with COVID-19.<sup>11, 12</sup> However, the putative benefits of supplementary vitamin D<sub>3</sub> to patients with COVID-19 remain speculative and partially supported by limited data from observational studies and one small-scale, non-randomized clinical trial.<sup>13-15</sup> To our knowledge, this is the first randomized, double-blind, placebo-controlled trial to investigate the safety and efficacy of vitamin D<sub>3</sub> supplementation on hospital length of stay and other relevant clinical outcomes in hospitalized patients with severe COVID-19. Our main *a priori* hypothesis was that a single dose of 200,000 IU of vitamin D<sub>3</sub> supplementation would increase 25-hydroxyvitamin D levels and shorten hospital length of stay among these patients.

# **Methods**

The study was approved by the Ethics Committee of Clinical Hospital of the School of Medicine of the University of Sao Paulo and by the Ethics Committee of Ibirapuera Field Hospital. All the procedures were conducted in accordance with the Declaration of Helsinki. The participants provided written informed consent before being enrolled in the study (Ethics Committee Approval Number 30959620.4.0000.0068). The trial protocol and statistical plan are included in Supplement 1. This manuscript was written according to the recommendations by the Consolidated Standards of Reporting Trials (CONSORT) guidelines (see Supplement 2).

# **Participants**

Hospitalized patients were recruited from Clinical Hospital of the School of Medicine of the University of Sao Paulo (a quaternary referral teaching hospital), and from Ibirapuera Field Hospital, both located in Sao Paulo, Brazil. Enrollment started on June 2, 2020, to August 27, 2020, with the final follow-up on October 7, 2020.

# **Inclusion criteria**

Inclusion criteria were: 1) adults aged 18 years or older; 2) diagnosis of COVID-19 by either polymerase chain reaction (PCR) for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from nasopharyngeal swabs or computed tomography scan findings (bilateral multifocal ground-glass opacities  $\geq$  50%) compatible with the disease; 3) diagnosis of flu syndrome with hospitalization criteria on hospital admission, presenting respiratory rate  $\geq$  24 breaths per minute, saturation  $<$  93% on room air or risk factors for complications, such as heart disease, diabetes

mellitus, systemic arterial hypertension, neoplasms, immunosuppression, pulmonary tuberculosis, and obesity, followed by COVID-19 confirmation before randomization.

#### **Exclusion criteria**

Exclusion criteria were: 1) patient unable to read and sign the written informed consent; 2) patient already admitted under invasive mechanical ventilation; 3) previous vitamin D<sub>3</sub> supplementation (> 1000 IU/day); 4) renal failure requiring dialysis or creatinine  $\geq$  2.0 mg/dL; 5) hypercalcemia defined by total calcium > 10.5 mg/dL; 6) pregnant or lactating women; and 7) patients with expected hospital discharge in less than 24 hours.

#### **Study design and treatment**

This was a multicenter, double-blind, parallel-group, randomized placebo-controlled trial. Eligibility screening was performed between June 2, 2020 to July 21, 2020 at Clinical Hospital of the School of Medicine of the University of Sao Paulo (n = 122), and from July 22, 2020 to August 27, 2020 at Ibirapuera Field Hospital (n = 118). The final follow-up in both centers was on October 7, 2020. Eligible patients were assigned in a 1:1 ratio into either the vitamin D<sub>3</sub> supplementation group or the placebo group. The randomization list was created using a computer-generated code, which was managed by a staff member who had no role in the study. We assessed patients' clinical status, coexisting chronic diseases, demographic characteristics, self-reported body weight and height, and ethnicity on hospital admission. Outcomes were assessed at baseline and on hospital discharge or death records.

The vitamin D<sub>3</sub> supplementation group received an oral, single dose of 200,000 IU of vitamin D<sub>3</sub> dissolved in a 10 mL of peanut oil solution on the same day of randomization. The selected dose is within the recommended range for effectively

promoting vitamin D sufficiency.<sup>16</sup> Patients in the placebo group received 10 mL of peanut oil solution. The vitamin D<sub>3</sub> and placebo solutions were identical in color, taste, smell, consistency, and container. Both were prepared by the pharmacy unit of Clinical Hospital and labeled by a staff member who did not participate in the study. Allocation blindness was kept until the final statistical analysis.

## **Outcome measures**

The primary outcome was hospital length of stay, defined as the total number of days that patients remained hospitalized from the date of study admission until the date of hospital discharge or death. The criteria used for patient discharge were: 1) no need for supplemental oxygen in the last 48 hours; 2) no fever in the last 72 hours; and 3) oxygen saturation > 93% in room air without respiratory distress.

The secondary outcomes were: 1) mortality; 2) number of patients admitted to the intensive care unit (ICU); 3) number of patients who needed mechanical ventilation and duration of mechanical ventilation; and 4) serum levels of 25-hydroxyvitamin D (assessed by a chemiluminescent immunoassay), calcium (assessed by a NM-BAPTA method), creatinine (assessed by a colorimetric assay based on kinetic Jaffe's reaction), and C-reactive protein and D-dimer (both assessed by an immunoturbidimetric assay).

The biochemical analyses were carried out in an accredited laboratory from Clinical Hospital.

## **Statistical Analysis**

Considering the lack of data available for sample size determination based on the primary outcome (i.e., hospital length of stay after vitamin D<sub>3</sub> supplementation in patients with severe COVID-19), the number of participants was chosen on the basis of



feasibility, such as resources, capacity of research staff and facility, and available patients, in line with current recommendations.<sup>17, 18</sup> Subsequently, we calculated sample size assuming a 50% between-group difference in hospital length of stay (considering 7 days as a median time of stay, with an expected variability of 9 days). By considering a power of 80% and a 2-sided significance level of 5% ( $\alpha = .05$ ), the total sample was estimated to be 208 patients (104 in each arm). Considering possible dropouts, and to increase the power for secondary outcomes, we opted by increasing the sample size by approximately 15%.

All analyses were carried out following the intention-to-treat principle for all randomized patients, with no imputation for any missing data. Proportions were compared between groups using  $\chi^2$  test and Fisher's exact test. Student's t-tests were used for comparing continuous variables at baseline. The log-rank test was used to compare the Kaplan-Meier estimate curves the number of days for hospital length of stay, the primary outcome. Cox regression models for hospital length of stay, admission to ICU and mechanical ventilation requirement were adjusted by potential confounders that were not fully balanced by randomization ( $P < .2$ ) to estimate hazard ratios (HR), with corresponding 2-sided 95% CI. Generalized estimating equations (GEE) for repeated measures were used for testing possible differences in laboratory parameters, assuming group and time as fixed factors, with marginal distribution, and a first-order autoregressive correlation matrix to test the main and interaction effects. *Post-hoc* tests with Bonferroni's adjustment were performed for multiple comparisons. The aforementioned statistical procedures were also carried out in *post-hoc* sensitivity analyses involving patients exhibiting 25-hydroxyvitamin D deficiency (i.e.,  $< 20$  ng/mL).

- 1 Statistical analyses were performed with IBM-SPSS software, version 20.0.
- 2 Significance level was set at  $\alpha = .05$ .

# Results

## Patients

Of 1208 patients assessed for eligibility, 240 were eligible and randomly assigned to either the vitamin D<sub>3</sub> group or the placebo group. Patients were non-eligible due to the following reasons: 284 were at ICU, 263 had hospital discharge within 24 hours, 217 did not have COVID-19 confirmation, 95 had renal dysfunction, 37 had dementia or severe mental confusion hampering their ability to provide the informed consent for participation, 30 were pregnant or lactating women, 14 had hypercalcemia due to metastatic neoplasm, 11 were receiving vitamin D<sub>3</sub> ( $\geq 1000$  IU/day), 9 were younger than 18 years, 6 were illiterate and, therefore, unable to read and sign the informed consent, and 2 died before randomization.

Of the 120 patients who were randomized to the vitamin D<sub>3</sub> group, 3 did not receive intervention (1 withdrew the consent, 1 vomited immediately after ingesting the supplement, and 1 was admitted to the ICU before taking vitamin D<sub>3</sub>) and 3 were lost to follow-up. Of the 120 patients who were randomized to the placebo group, 2 withdrew the consent. Thus, of the 240 patients randomized, 232 (96.7%) completed the follow-up (**Figure 1**).

Overall, patients' age was 56.3 years (SD, 14.6), BMI was 31.6 kg/m<sup>2</sup> (SD, 7.1), 56.3% were men, 55% were white, 52.5% had hypertension, 35% had diabetes, 13.3% had cardiovascular diseases, and 6.3% had asthma. The mean time between the onset of symptoms and randomization was 10.2 days (SD, 4.3); 89.6% required supplemental oxygen at baseline (183 were on oxygen therapy and 32 were on non-invasive ventilation), and 59.6% had computed tomography scan findings suggestive of COVID-19. Demographic and clinical characteristics did not significantly differ between groups, except for sore throat, which was more prevalent in the vitamin D<sub>3</sub> group vs placebo

(38.3% vs 24.2%,  $P = .026$ ), and PTH, which was higher in the vitamin D<sub>3</sub> group vs placebo (50.1 vs 42.6 pg/mL,  $P = .025$ ) (**Table 1**).

## Primary Outcome

Hospital length of stay (**Figure 2**) was comparable between the vitamin D<sub>3</sub> group and the placebo group (7.0 days [95% CI, 6.1 to 7.9] and 7.0 days [95% CI, 6.2 to 7.8 days], HR, 1.12, [95% CI, 0.9 to 1.5];  $P = .379$ ; respectively). The Cox regression model did not show any significant associations between this outcome and potential confounders.

## Secondary Outcomes

There were no significant differences between the vitamin D<sub>3</sub> group and the placebo group for mortality (7.0% vs 5.1%;  $P = .590$ ), admission to ICU (15.8% vs 21.2%;  $P = .314$ ) and mechanical ventilation requirement (7.0% vs 14.4%;  $P = .090$ ) (**Figure 3**).

Duration of mechanical ventilation was also comparable between the vitamin D<sub>3</sub> group (18.1 days [95% CI, 3.5 to 32.7]) and the placebo group (11.4 days [95% CI, 7.1 to 15.6];  $P = .549$ , respectively).

The Cox regression model did not show significant associations between secondary outcomes and potential confounders.

Vitamin D<sub>3</sub> supplementation significantly increased 25-hydroxyvitamin D levels vs placebo (difference, 24.0 ng/mL [95% CI, 21.1- 26.9];  $P < .001$ ) (**Figure 3**). Following the intervention, 86.7% of the patients in the vitamin D<sub>3</sub> group showed 25-hydroxyvitamin D levels above 30 ng/mL (vs 10.9% in the placebo group), and only 6.7% of the patients in the vitamin D<sub>3</sub> group exhibited 25-hydroxyvitamin D deficiency (vs 51.5% in the placebo group).

## **Post-hoc Sensitivity Analyses**

In a sensitivity analysis involving patients with 25-hydroxyvitamin D deficiency at baseline (n = 116) (**Supplementary Table 1**), vitamin D<sub>3</sub> supplementation significantly increased 25-hydroxyvitamin D levels vs placebo (difference, 22.7 ng/mL [95% CI, 19.3 to 26.1];  $P < .001$ ) (**Figure 3**). Among the patients with 25-hydroxyvitamin D deficiency, no between-group differences were observed in length of hospital stay (**Figure 2**). In addition, there were no significant differences between the vitamin D<sub>3</sub> group and the placebo group for mortality (7.0% vs 1.7%;  $P = .206$ ), admission to ICU (17.5% vs 15.5%;  $P = .806$ ), and mechanical ventilation requirement (7.0% vs 8.6%;  $P > .999$ ) (**Figure 3**). Duration of mechanical ventilation did not differ between the vitamin D<sub>3</sub> group (15.0 days [95% CI, -12.0 to 42.0]) and the placebo group (12.6 days [95% CI, -7.6 to 26.0];  $P = .730$ ).

## **Safety and Adverse Events**

There were no changes in any health-related laboratory markers following the intervention (**Table 2**). Vitamin D<sub>3</sub> supplementation was well tolerated and no severe adverse events were reported throughout the trial, with the exception of one patient who vomited following vitamin D<sub>3</sub> administration.

## **Discussion**

This is the first randomized, double-blind, placebo-controlled trial to show that vitamin D<sub>3</sub> supplementation is safe and increases 25-hydroxyvitamin D levels, but is ineffective to improve hospital length of stay or any other clinical outcomes among hospitalized patients with severe COVID-19.

Vitamin D has been postulated to play an important role on immune system, acting as a regulator of both innate and adaptative responses.<sup>6, 19</sup> Observational studies have shown that 25-hydroxyvitamin D levels are associated with better clinical outcomes in respiratory diseases.<sup>20</sup> Positive associations between low 25-hydroxyvitamin D levels and poor prognosis among patients with COVID-19 have also been observed.<sup>21</sup> Furthermore, a small-scale, non-randomized trial demonstrated that the administration of regular boluses of vitamin D<sub>3</sub> before the infection was associated with better survival and less severe disease among older, frail patients with COVID-19.<sup>22</sup> In the current trial, however, a single dose of 200,000 IU of vitamin D<sub>3</sub> supplementation failed to promote any clinically relevant effects among hospitalized patients with severe COVID-19, contesting the utility of supplementary vitamin D<sub>3</sub> as a treatment in this disease. The lack of clinical benefits seen in this study was independent of the ability of vitamin D<sub>3</sub> supplementation to increase serum 25-hydroxyvitamin D levels. In fact, following the intervention, 86.7% of the patients in the supplementation arm achieved vitamin D sufficiency ( $\geq 30$  ng/mL) vs 11% only in the placebo group. In a sensitivity analysis confined to the patients exhibiting 25-hydroxyvitamin D deficiency, vitamin D<sub>3</sub> supplementation remained effective in increasing 25-hydroxyvitamin D levels vs placebo; yet, no clinical improvements were noted. Collectively, these analyses indicate that a single oral dose of 200,000 IU of supplementation can rapidly increase 25-hydroxyvitamin levels, in agreement with our hypothesis, so that the present null findings cannot be attributed to the failure of increasing serum 25-hydroxyvitamin D levels. Despite the clinical inefficacy of vitamin D<sub>3</sub> supplementation, the intervention was not associated with any important adverse events or meaningful changes in laboratory

parameters, suggesting that a relatively high-dose of vitamin D<sub>3</sub> can be well tolerated in general and free of adverse effects in patients with COVID-19. The strengths of this study include the randomized, double-blind, placebo-controlled experimental design, the adequate power, particularly for the primary analysis, the very low attrition rate (3.3%), the concomitant assessment of 25-hydroxyvitamin D levels along with clinical outcomes, and the assessment of hospitalized patients with severe COVID-19.

## Limitations

This trial has several limitations. First, the sample size could have been underpowered to detect significant changes for the secondary outcomes. Second, as the patients had several coexisting diseases and were subjected to a diverse medication regimen, the results could have been affected by the heterogeneity of the sample and its treatment. Third, the proportion of patients with 25-hydroxyvitamin D deficiency enrolled in this study was considerably lower than those reported in other cohorts,<sup>23</sup> possibly as a consequence of differences in geographic locations. Although we conducted sensitivity analyses involving patients with 25-hydroxyvitamin D deficiency, one could argue that they could have been underpowered, as previously pointed out. Therefore, caution should be exercised in generalizing these findings to patients from other geographical regions. Finally, the findings should be also confined to the dose and supplementation strategy used in this trial. Further studies should determine whether preventive or early vitamin D<sub>3</sub> supplementation could be useful in the treatment of patients with COVID-19, especially those with a mild or moderate disease.

## 1    **Conclusions**

2    Among hospitalized patients with severe COVID-19, a single dose of 200,000 IU of  
3    vitamin D<sub>3</sub> supplementation was safe and increased 25-hydroxyvitamin D levels, but  
4    did not reduce hospital length of stay or any other clinically relevant outcomes vs  
5    placebo. Thus, this trial does not support the use of vitamin D<sub>3</sub> supplementation as an  
6    adjuvant treatment of patients with COVID-19.



# 1 Article information

2 **Author Affiliations:** Rheumatology Division, Hospital das Clinicas HCFMUSP,  
3 Faculdade de Medicina da Universidade de Sao Paulo, Sao Paulo, Brazil (Murai,  
4 Fernandes, Sales, Duran, Silva, Franco, Macedo, Dalmolin, Baggio, Balbi, Reis,  
5 Caparbo, Pereira); Applied Physiology & Nutrition Research Group; Faculdade de  
6 Medicina da Universidade de Sao Paulo, Sao Paulo, Brazil (Pinto, Goessler, Gualano);  
7 Clinical Pathology Division, Hospital das Clinicas HCFMUSP, Faculdade de Medicina  
8 da Universidade de Sao Paulo, Sao Paulo, Brazil (Antonangelo); Food Research Center,  
9 Universidade de Sao Paulo, Sao Paulo, Brazil (Gualano).

10 **Author Contributions:** Dr. Pereira had full access to all of the data in the study and  
11 take responsibility for the integrity of the data and the accuracy of the data analysis.

12 *Concept and design:* Murai, Fernandes, Pinto, Goessler, Gualano, Pereira.

13 *Acquisition, analysis and interpretation:* All authors.

14 *Drafting of the manuscript:* Murai, Fernandes, Gualano, Pereira.

15 *Critical revision of the manuscript for important intellectual content:* All authors.

16 *Statistical analysis:* Murai, Fernandes, Pinto, Reis, Gualano, Pereira.

17 *Obtained funding:* Gualano, Pereira.

18 *Supervision:* Gualano, Pereira.

19 *Administrative, technical, or material support:* Sales, Antonangelo, Caparbo.

20 **Conflict of Interest Disclosures:** The authors have nothing to disclose.

21 **Funding/Support:** This study was supported by Sao Paulo Research Foundation  
22 (FAPESP) (grants 20/05752-4; 19/24782-4; 20/11102-2; 16/00006-7; 17/13552-2;  
23 15/26937-4; 19/18039-7) and Conselho Nacional de Desenvolvimento Científico e  
24 Tecnológico (305556/2017-7).

- 1    **Data Sharing Statement:** See Supplement 3.
- 2    **Additional Contributions:** The authors are thankful to Dr. Monica Pinheiro and Dr.
- 3    Roberta Costa for the assistance at Ibirapuera Field Hospital; Dr. Rogério Ruscitto do
- 4    Prado for conducting statistical analyses; Mayara Diniz Santos for the technical support;
- 5    all the staff members from both centers; all the patients who participated in this study.

## References

1. Liu PT, Stenger S, Li H, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science*. Mar 2006;311(5768):1770-3. doi:10.1126/science.1123933
2. Aglipay M, Birken CS, Parkin PC, et al. Effect of High-Dose vs Standard-Dose Wintertime Vitamin D Supplementation on Viral Upper Respiratory Tract Infections in Young Healthy Children. *JAMA*. Jul 18 2017;318(3):245-254. doi:10.1001/jama.2017.8708
3. Campbell GR, Spector SA. Autophagy induction by vitamin D inhibits both Mycobacterium tuberculosis and human immunodeficiency virus type 1. *Autophagy*. Oct 2012;8(10):1523-5. doi:10.4161/auto.21154
4. van Etten E, Mathieu C. Immunoregulation by 1,25-dihydroxyvitamin D3: basic concepts. *J Steroid Biochem Mol Biol*. Oct 2005;97(1-2):93-101. doi:10.1016/j.jsbmb.2005.06.002
5. Laplana M, Royo JL, Fibla J. Vitamin D Receptor polymorphisms and risk of enveloped virus infection: A meta-analysis. *Gene*. Dec 2018;678:384-394. doi:10.1016/j.gene.2018.08.017
6. Bilezikian JP, Bikle D, Hewison M, et al. MECHANISMS IN ENDOCRINOLOGY: Vitamin D and COVID-19. *Eur J Endocrinol*. Nov 2020;183(5):R133-R147. doi:10.1530/EJE-20-0665
7. Autier P, Boniol M, Pizot C, Mullie P. Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol*. Jan 2014;2(1):76-89. doi:10.1016/S2213-8587(13)70165-7
8. Aibana O, Huang CC, Aboud S, et al. Vitamin D status and risk of incident tuberculosis disease: A nested case-control study, systematic review, and individual-participant data meta-analysis. *PLoS Med*. 09 2019;16(9):e1002907. doi:10.1371/journal.pmed.1002907
9. Martineau AR, Jolliffe DA, Hooper RL, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ*. Feb 2017;356:i6583. doi:10.1136/bmj.i6583
10. Sabetta JR, DePetrillo P, Cipriani RJ, Smardin J, Burns LA, Landry ML. Serum 25-hydroxyvitamin d and the incidence of acute viral respiratory tract infections in healthy adults. *PLoS One*. Jun 2010;5(6):e11088. doi:10.1371/journal.pone.0011088
11. Mitchell F. Vitamin-D and COVID-19: do deficient risk a poorer outcome? *Lancet Diabetes Endocrinol*. 07 2020;8(7):570. doi:10.1016/S2213-8587(20)30183-2
12. Martineau AR, Forouhi NG. Vitamin D for COVID-19: a case to answer? *Lancet Diabetes Endocrinol*. Sep 2020;8(9):735-736. doi:10.1016/S2213-8587(20)30268-0
13. Meltzer DO, Best TJ, Zhang H, Vokes T, Arora V, Solway J. Association of Vitamin D Status and Other Clinical Characteristics With COVID-19 Test Results. *JAMA Netw Open*. 09 2020;3(9):e2019722. doi:10.1001/jamanetworkopen.2020.19722
14. Kaufman HW, Niles JK, Kroll MH, Bi C, Holick MF. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. *PLoS One*. 2020;15(9):e0239252. doi:10.1371/journal.pone.0239252
15. Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res*. Jul 2020;32(7):1195-1198. doi:10.1007/s40520-020-01570-8

16. Kearns MD, Alvarez JA, Tangpricha V. Large, single-dose, oral vitamin D supplementation in adult populations: a systematic review. *Endocr Pract.* Apr 2014;20(4):341-51. doi:10.4158/EP13265.RA
17. Bacchetti P. Current sample size conventions: flaws, harms, and alternatives. *BMC Med.* Mar 2010;8:17. doi:10.1186/1741-7015-8-17
18. Bacchetti P, McCulloch CE, Segal MR. Simple, defensible sample sizes based on cost efficiency. *Biometrics.* Jun 2008;64(2):577-85; discussion 586-94. doi:10.1111/j.1541-0420.2008.01004\_1.x
19. Franco AS, Freitas TQ, Bernardo WM, Pereira RMR. Vitamin D supplementation and disease activity in patients with immune-mediated rheumatic diseases: A systematic review and meta-analysis. *Medicine (Baltimore).* Jun 2017;96(23):e7024. doi:10.1097/MD.00000000000007024
20. Cannell JJ, Vieth R, Umhau JC, et al. Epidemic influenza and vitamin D. *Epidemiol Infect.* Dec 2006;134(6):1129-40. doi:10.1017/S0950268806007175
21. Carpagnano GE, Di Lecce V, Quaranta VN, et al. Vitamin D deficiency as a predictor of poor prognosis in patients with acute respiratory failure due to COVID-19. *J Endocrinol Invest.* Aug 2020;doi:10.1007/s40618-020-01370-x
22. Annweiler G, Corvaisier M, Gautier J, et al. Vitamin D Supplementation Associated to Better Survival in Hospitalized Frail Elderly COVID-19 Patients: The GERIA-COVID Quasi-Experimental Study. *Nutrients.* Nov 2020;12(11)doi:10.3390/nu12113377
23. Hernández JL, Nan D, Fernandez-Ayala M, et al. Vitamin D Status in Hospitalized Patients With SARS-CoV-2 Infection. *The Journal of Clinical Endocrinology & Metabolism.* 2020;doi:10.1210/clinem/dgaa733

## Figure legends

### Figure 1. Flow of patients.

**Figure 2. Kaplan-Meier curves for hospital length of stay.** Vertical bars present single censored events. The shaded areas represent the 95% confidence intervals. The adjusted hazard ratio for total number of days that patients remained hospitalized from the date of study admission until the date of hospital discharge or death was 1.12 (95% CI, 0.9 to 1.5;  $P = .379$ ) for the vitamin D<sub>3</sub> group (7.0 days [95% CI, 6.1 to 7.9]) vs the placebo group (7.0 days [95% CI, 6.2 to 7.8 days]).


**Figure 3. Serum 25-hydroxyvitamin D levels, mortality, admission to intensive care unit (ICU), and mechanical ventilation requirement.** The panels show the comparisons between the vitamin D<sub>3</sub> and placebo group for all patients ( $n = 240$ ) (Panels A and B) and for those with 25-hydroxyvitamin D deficiency ( $< 20$  ng/mL) ( $n = 116$ ) (Panels C and D). For all patients, no significant differences were found between the vitamin D<sub>3</sub> group and the placebo group for mortality (7.0% vs 5.1%;  $P = .590$ ), admission to ICU (15.8% vs 21.2%;  $P = .314$ ), and need of mechanical ventilation (7.0% vs 14.4%;  $P = .090$ ). Vitamin D<sub>3</sub> supplementation significantly increased 25-hydroxyvitamin D levels vs placebo (difference, 24.0 ng/mL [95% CI, 21.1- 26.9];  $P < .001$ ). For patients with 25-hydroxyvitamin D deficiency, there were no significant differences between the vitamin D<sub>3</sub> group and the placebo group for mortality (7.0% vs 1.7%;  $P = .206$ ), admission to ICU (17.5% vs 15.5%;  $P = .806$ ), and mechanical ventilation requirement (7.0% vs 8.6%;  $P > .999$ ). Vitamin D<sub>3</sub> supplementation significantly increased 25-hydroxyvitamin D levels vs placebo (difference, 22.7 ng/mL

[95% CI, 19.3- 26.1];  $P < .001$ ). Box plots depict median and interquartile range. Outliers (i.e., defined as a value  $< 5$  or  $> 95$  percentiles) were represented by filled circles. \* means  $P < .05$  between Baseline and Post; # means  $P < .05$  between groups at Post.

**Table 1. Baseline demographic and clinical characteristics.**


**Table 2. Laboratory variables.**

**Supplementary Table 1. Baseline demographic and clinical characteristics from patients with 25-hydroxyvitamin D deficiency (< 20 ng/mL).**




**Figure 1. Flow of patients.**





**Figure 2. Kaplan-Meier curves for hospital length of stay.**



**Figure 3. Serum 25-hydroxyvitamin D levels, mortality, admission to intensive care unit (ICU), and mechanical ventilation requirement.** \* means  $P < .05$  between Baseline and Post; # means  $P < .05$  between groups at Post.

**Table 1. Baseline Demographic and Clinical Characteristics**

	Vitamin D <sub>3</sub> (n = 120)	Placebo (n = 120)	<i>P</i> value
Age, mean (SD), y	56.8 (14.2)	55.8 (15.0)	.584
Sex, No. (%)			
Male	70 (58.3)	65 (54.2)	.515
Female	50 (41.7)	55 (45.8)	
Race, No. (%)			
White	62 (51.7)	70 (58.3)	.399
Brown	37 (30.8)	36 (30.0)	
Black	20 (16.7)	14 (11.7)	
Asian	1 (0.8)	0 (0.0)	
Days since symptoms onset, mean (SD)	10.3 (4.7) [n=116]	10.2 (3.8) [n=119]	.787
Body mass index, mean (SD), kg/m <sup>2</sup>	31.9 (6.5) [n=109]	31.3 (7.6) [n=110]	.548
Underweight, No./total (%)	0 (0)	2 (1.8)	.067
Normal, No./total (%)	9/109 (8.3)	19/110 (17.3)	
Overweight, No./total (%)	37/109 (33.9)	31/110 (28.2)	
Obesity, No./total (%)	63/109 (57.8)	58/110 (52.7)	
Acute COVID-19 symptoms, No. (%)			
Fever	86 (71.7)	81 (67.5)	.575
Cough	103 (85.8)	99 (82.5)	.596
Fatigue	98 (81.7)	100 (83.3)	.865
Joint pain	46 (38.3)	35 (29.2)	.172
Myalgia	69 (57.5)	71 (59.2)	.896
Nasal congestion	39 (32.5)	43 (35.8)	.683
Runny nose	44 (36.7)	44 (36.7)	>.999
Sore throat	46 (38.3)	29 (24.2)	.026
Diarrhea	41 (34.2)	46 (38.3)	.591
Coexisting diseases, No. (%)			
Hypertension	68 (56.7)	58 (48.3)	.196
Cardiovascular disease	16 (13.3)	16 (13.3)	>.999
Diabetes	49 (40.8)	35 (29.2)	.058
Chronic obstructive pulmonary disease	7 (5.8)	5 (4.2)	.554
Asthma	8 (6.7)	7 (5.8)	.790

Chronic kidney disease	2 (1.7)	0 (0.0)	.489
Rheumatic disease	13 (10.8)	10 (8.3)	.511
Concomitant medications, No. (%)			
Antibiotic	102 (85.0)	105 (87.5)	.708
Anticoagulant	110 (91.7)	103 (85.8)	.220
Analgesic	45 (37.5)	52 (43.7)	.430
Corticosteroids	77 (64.2)	73 (60.8)	.689
Antihypertensive	67 (55.8)	56 (46.7)	.196
Hypoglycemic	26 (21.7)	24 (20.0)	.874
Hypolipidemic	15 (12.5)	18 (15.0)	.708
Antiemetic	45 (37.5)	55 (45.8)	.239
Antiviral	4 (3.3)	4 (3.3)	>.999
Proton pump inhibitor	47 (39.2)	49 (40.8)	.895
Thyroid	10 (8.3)	10 (8.3)	>.999
Oxygen supplementation, No. (%)			
No oxygen therapy	16 (13.3)	9 (7.5)	
Oxygen therapy	86 (71.7)	97 (80.8)	.210
Non-invasive ventilation	18 (15.0)	14 (11.7)	
Computed tomography findings, No. (%)			
Ground-glass opacities < 50%	61 (50.8)	66 (55.0)	
Ground-glass opacities ≥ 50%	47 (39.2)	39 (32.5)	.543
Not available	12 (10.0)	15 (12.5)	
Laboratory variables			
Haemoglobin, mean (SD), g/L	13.1 (2.1)	12.8 (2.1)	.298
Neutrophils count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	6.7 (4.0) [n = 119]	7.2 (3.6) [n = 120]	.281
Lymphocyte count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	1.2 (0.5) [n = 119]	1.1 (0.8) [n = 120]	.849
Platelet count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	305.4 (116.4)	286.7 (128.5)	.239
Erythrocyte sedimentation rate, mean (SD), mm	58.4 (40.6) [n = 117]	60.9 (36.7) [n = 119]	.627
C-reactive protein, mean (SD), mg/L	79.6 (75.2) [n = 119]	90.4 (80.4) [n = 120]	.286
D-dimer, mean (SD), ng/mL	2091 (5001.3) [n = 119]	1720.7 (3630.1) [n = 119]	.514
Albumin, mean (SD), g/L	3.1 (0.5) [n = 110]	3.0 (0.4) [n = 100]	.454
Gamma globulins, mean (SD), g/L	1.1 (0.4) [n = 110]	1.1 (0.3) [n = 100]	.337

Creatinine, mean (SD), mg/dL	0.9 (0.3)	0.8 (0.2)	.149
Urea, mean (SD), mg/dL	40.1 (20.1) [n = 120]	37.8 (14.5) [n = 119]	.309
Phosphorus, mean (SD), mg/dL	3.0 (0.6) [n = 117]	3.0 (0.8) [n = 116]	.790
25-hydroxyvitamin D, mean (SD), ng/mL	21.0 (10.2) [n = 118]	20.6 (8.1) [n = 118]	.747
Parathyroid hormone, mean (SD), pg/mL	50.1 (27.3) [n = 113]	42.6 (21.5) [n = 110]	.025
Calcium, total, mean (SD), mg/dL	8.7 (0.5) [n = 118]	8.7 (0.5) [n = 119]	.811
Total cholesterol, mean (SD), mg/dL	164.3 (43.3) [n = 116]	164.5 (47.2) [n = 115]	.975
LDL-cholesterol, mean (SD), mg/dL	101.3 (34.6) [n = 116]	99.3 (39.1) [n = 114]	.681
HDL-cholesterol, mean (SD), mg/dL	34.6 (11.4) [n = 116]	34.5 (11.0) [n = 114]	.916
Triglycerides, mean (SD), mg/dL	178.6 (75.5) [n=116]	192.5 (93.6) [n = 114]	.218

For continuous variables, groups were compared using independent t-test.

For categorical variables, groups were compared using  $\chi^2$  test or Fisher's exact test, as appropriate.

**Table 2. Laboratory variables.**

All patients	Vitamin D <sub>3</sub> group		Difference	Placebo group		Difference	<i>P</i> value <sup>a</sup>	<i>P</i> value <sup>b</sup>
	Baseline	Post	(Baseline – Post)	Baseline	Post	(Baseline – Post)		
Variables								
Haemoglobin, mean (SD), g/L	13.1 (2.1)	12.7 (2.4)	0.4 (3.0) [n = 111]	12.8 (2.1)	12.7 (2.1)	0.2 (3.1) [n = 114]	.525	.595
Neutrophils count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	6.7 (4.0)	7.0 (4.2)	-0.4 (3.6) [n = 111]	7.2 (3.6)	7.1 (4.9)	0.1 (5.0) [n = 114]	.460	.426
Lymphocyte count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	1.2 (0.5)	1.8 (0.9)	-0.7 (0.8) [n = 111]	1.1 (0.8)	2.0 (1.1)	-0.9 (0.8) [n = 114]	.060	.075
Platelet count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	305.4 (116.4)	367.5 (137.9)	-63.3 (100.7) [n = 111]	286.7 (128.5)	357.7 (141.1)	-70.6 (106.7) [n = 114]	.566	.597
Erythrocyte sedimentation rate, mean (SD), mm	58.4 (40.6)	48.6 (39.1)	8.9 (33.0) [n = 101]	60.9 (36.7)	44.2 (33.6)	17.1 (36.1) [n = 99]	.110	.102
C-reactive protein, mean (SD), mg/L	79.6 (75.2)	28.2 (54.0)	47.9 (70.6) [n = 109]	90.4 (80.4)	22.9 (42.1)	61.3 (79.1) [n = 107]	.156	.190
D-dimer, mean (SD), ng/mL	2091 (5001.3)	1589.4 (2484.8)	592.6 (5150.7) [n = 119]	1720.7 (3630.1)	1284.0 (1791.4)	316.3 (2049.7) [n = 119]	.843	.620
Creatinine, mean (SD), mg/dL	0.9 (0.3)	1.0 (0.7)	-0.1 (0.6) [n = 112]	0.8 (0.2)	0.9 (0.6)	-0.1 (0.7) [n = 112]	.928	.927
Urea, mean (SD), mg/dL	40.1 (20.1)	49.0 (46.8)	-9.6 (42.4) [n = 112]	37.8 (14.5)	46.6 (50.4)	-8.4 (49.4) [n = 113]	.956	.839
Phosphorus,	3.0 (0.6)	3.5 (0.7)	-0.5 (0.8)	3.0 (0.8)	3.7 (1.3)	-0.6 (1.4)	.334	.596

mean (SD), mg/dL			[n = 108]			[n = 108]		
Parathyroid hormone, mean (SD), pg/mL	50.1 (27.3)	35.0 (19.8)	13.3 (23.3) [n = 100]	42.6 (21.5)	36.7 (19.5)	7.5 (16.1) [n = 99]	.013	.049
Total calcium, mean (SD), mg/dL	8.7 (0.5)	9.1 (0.6)	-0.4 (0.5) [n = 106]	8.7 (0.5)	9.1 (0.5)	-0.4 (0.6) [n = 106]	.968	.890
Total cholesterol, mean (SD), mg/dL	164.3 (43.3)	183.2 (48.7)	-16.0 (30.4) [n = 104]	164.5 (47.2)	187.1 (50.3)	-22.5 (35.8) [n = 101]	.137	.170
LDL-cholesterol, mean (SD), mg/dL	101.3 (34.6)	111.9 (37.0)	-8.0 (24.8) [n = 104]	99.3 (39.1)	112.0 (40.2)	-12.5 (28.1) [n = 101]	.223	.243
HDL-cholesterol, mean (SD), mg/dL	34.6 (11.4)	37.5 (11.1)	-2.9 (8.0) [n = 104]	34.5 (11.0)	37.5 (10.3)	-3.7 (11.0) [n = 101]	.666	.553
Triglycerides, mean (SD), mg/dL	178.6 (75.5)	221.7 (115.3)	-41.1 (88.7) [n = 104]	192.5 (93.6)	251.9 (129.2)	-54.5 (106.6) [n = 101]	.244	.336

<sup>a</sup> *P* value represents time by group interaction, calculated by Generalized Estimating Equations (GEE) with normal distribution and identity link function with AR (1) correlation matrix.

<sup>b</sup> *P* value represents between-group comparisons for the difference, calculated by independent t-test.

**Supplementary Table 1. Baseline Demographic and Clinical Characteristics from patients with 25-hydroxyvitamin D deficiency (< 20 ng/mL).**

	Vitamin D <sub>3</sub> (n = 58)	Placebo (n = 58)	<i>P</i> value
Age, mean (SD), y	58.9 (15.4)	56.1 (14.3)	.308
Sex, No. (%)			
Male	31 (53.4)	28 (48.3)	.577
Female	27 (46.6)	30 (51.7)	
Race, No. (%)			
White	26 (44.8)	34 (58.6)	.340
Brown	19 (32.8)	15 (25.9)	
Black	12 (20.7)	9 (15.5)	
Asian	1 (1.7)	0 (0)	
Days since symptoms onset, mean (SD)	10.5 (5.7)	10.4 (3.7)	.932
Body mass index, mean (SD), kg/m <sup>2</sup>	32.0 (6.8) [n = 50]	31.7 (8.4) [n = 54]	.832
Underweight, No./total (%)	0 (0)	2/54 (3.7)	.275
Normal, No./total /(%)	5/50 (10.0)	8/54 (14.8)	
Overweight, No./total /(%)	15/50 (30.0)	18/54 (33.3)	
Obesity, No./total (%)	30/50 (60.0)	26/54 (48.1)	
Acute COVID-19 symptoms			
Fever	39 (67.2)	40 (69.0)	.842
Cough	50 (86.2)	51 (87.9)	.782
Fatigue	47 (81.0)	49 (84.5)	.623
Joint pain	20 (34.5)	15 (25.9)	.312
Myalgia	33 (56.9)	35 (60.3)	.706
Nasal congestion	16 (27.6)	26 (44.8)	.053
Runny nose	21 (36.2)	22 (37.9)	.848
Sore throat	24 (41.4)	13 (22.4)	.028
Diarrhea	16 (27.6)	22 (37.9)	.235
Coexisting diseases, No. (%)			
Hypertension	35 (60.3)	28 (48.3)	.192
Cardiovascular disease	6 (10.3)	8 (13.8)	.569
Diabetes	27 (46.6)	19 (32.8)	.129
Chronic obstructive pulmonary disease	5 (8.6)	5 (8.6)	>.999



Asthma	4 (6.9)	2 (3.4)	.679
Chronic kidney disease	1 (1.7)	0 (0.0)	>.999
Rheumatic disease	6 (10.3)	3 (5.2)	.490
Concomitant medications, No. (%)			
Antibiotic	47 (81.0)	50 (86.2)	.452
Anticoagulant	52 (89.7)	45 (77.6)	.079
Analgesic	23 (39.7)	30 (51.7)	.192
Corticosteroids	35 (60.3)	32 (55.2)	.573
Antihypertensive	34 (58.6)	29 (50.0)	.351
Hypoglycemic	12 (20.7)	13 (22.4)	.821
Hypolipidemic	9 (15.5)	9 (15.5)	>.999
Antiemetic	19 (32.8)	24 (41.4)	.336
Antiviral	2 (3.4)	2 (3.4)	>.999
Proton pump inhibitor	18 (31.0)	24 (41.4)	.246
Thyroid	6 (10.3)	5 (8.6)	.751
Oxygen supplementation, No. (%)			
No oxygen therapy	11 (19.0)	3 (5.2)	
Oxygen therapy	38 (65.5)	52 (89.7)	.008
Non-invasive ventilation	9 (15.5)	3 (5.2)	
Computed tomography findings, No. (%)			
Ground-glass opacities < 50%	27 (46.5)	18 (31.0)	
Ground-glass opacities ≥ 50%	24 (41.4)	31 (53.4)	.246
Not available	7 (12.1)	9 (15.5)	
Laboratory variables			
Haemoglobin, mean (SD), g/L	13.3 (2.2)	12.6 (2.1)	.114
Neutrophils count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	6.02 (3.73)	7.55 (3.82)	.031
Lymphocyte count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	1.19 (0.53)	1.10 (0.52)	.353
Platelet count, mean (SD), x10 <sup>3</sup> /mm <sup>3</sup>	301.8 (130.7)	292.3 (141.9)	.707
Erythrocyte sedimentation rate, mean (SD), mm	62.3 (39.3)	66.4 (40.0)	.587
C-reactive protein, mean (SD), mg/L	78.5 (72.6)	81.4 (82.7)	.843
D-dimer, mean (SD), ng/mL	2,950 (6,957)	2,044 (4,541)	.412
Albumin, mean (SD), g/L	2.95 (0.49)	2.98 (0.45)	.801

Gamma globulins, mean (SD), g/L	1.15 (0.44)	1.12 (0.33)	.665
Creatinine, mean (SD), mg/dL	0.86 (0.32)	0.84 (0.27)	.769
Urea, mean (SD), mg/dL	38.7 (19.2)	38.8 (14.9)	.975
Phosphorus, mean (SD), mg/dL	2.97 (0.64)	3.09 (0.73)	.356
25-hydroxyvitamin D, mean (SD), ng/mL	12.6 (4.0)	13.9 (4.7)	.108
Parathyroid hormone, mean (SD), pg/mL	52.7 (30.0)	43.9 (24.4)	.098
Calcium, total, mean (SD), mg/dL	8.54 (0.52)	8.67 (0.52)	.170
Total cholesterol, mean (SD), mg/dL	161.3 (46.1)	169.6 (50.8)	.369
LDL-cholesterol, mean (SD), mg/dL	98.7 (36.8)	104.3 (41.6)	.453
HDL-cholesterol, mean (SD), mg/dL	35.2 (13.0)	32.8 (12.2)	.318
Triglycerides, mean (SD), mg/dL	170.3 (80)	201.0 (91.9)	.062

---

For continuous variables, groups were compared using independent t-test.

For categorical variables, groups were compared using  $\chi^2$  test or Fisher's exact test, as appropriate.