



'Scientific Strabismus' or two related pandemics: coronavirus disease and vitamin D deficiency

Murat Kara¹, Timur Ekiz^{2*}, Vincenzo Ricci³, Özgür Kara⁴, Ke-Vin Chang⁵ and Levent Özçakar¹

¹Department of Physical Medicine and Rehabilitation, Hacettepe University Medical School, Ankara, Turkey

²Department of Physical Medicine and Rehabilitation, Türkmenbaşı Medical Center, Adana, Turkey

³Department of Biomedical and Neuromotor Science, Physical and Rehabilitation Medicine Unit, IRCCS Rizzoli Orthopaedic Institute, Bologna, Italy

⁴Geriatrics Unit, Yenimahalle Training and Research Hospital, Yıldırım Beyazıt University, Ankara, Turkey

⁵Department of Physical Medicine and Rehabilitation, National Taiwan University Hospital, Bei-Hu Branch, Taipei, Taiwan

(Submitted 5 April 2020 – Final revision received 4 May 2020 – Accepted 7 May 2020)

Abstract

The WHO has announced the novel coronavirus disease (COVID-19) outbreak to be a global pandemic. The distribution of community outbreaks shows seasonal patterns along certain latitude, temperature and humidity, that is, similar to the behaviour of seasonal viral respiratory tract infections. COVID-19 displays significant spread in northern mid-latitude countries with an average temperature of 5–11°C and low humidity. Vitamin D deficiency has also been described as pandemic, especially in Europe. Regardless of age, ethnicity and latitude, recent data showed that 40% of Europeans are vitamin D deficient (25-hydroxyvitamin D (25(OH)D) levels <50 nmol/l), and 13% are severely deficient (25(OH)D < 30 nmol/l). A quadratic relationship was found between the prevalences of vitamin D deficiency in most commonly affected countries by COVID-19 and the latitudes. Vitamin D deficiency is more common in the subtropical and mid-latitude countries than the tropical and high-latitude countries. The most commonly affected countries with severe vitamin D deficiency are from the subtropical (Saudi Arabia 46%; Qatar 46%; Iran 33.4%; Chile 26.4%) and mid-latitude (France 27.3%; Portugal 21.2%; Austria 19.3%) regions. Severe vitamin D deficiency was found to be nearly 0% in some high-latitude countries (e.g. Norway, Finland, Sweden, Denmark and Netherlands). Accordingly, we would like to call attention to the possible association between severe vitamin D deficiency and mortality pertaining to COVID-19. Given its rare side effects and relatively wide safety, prophylactic vitamin D supplementation and/or food fortification might reasonably serve as a very convenient adjunct therapy for these two worldwide public health problems alike.

Key words: Coronavirus: Death: Insufficiency: Europe: Acute respiratory syndrome

On 11 March 2020, the WHO announced the novel coronavirus disease (COVID-19) outbreak to be a global pandemic⁽¹⁾. The spread of COVID-19 is becoming unstoppable, and as of 15 May, more than 4 500 000 people have been infected and more than 300 000 people have died (Fig. 1)⁽¹⁾. The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the pathogen of COVID-19. SARS-CoV-2, classified into two β coronaviruses, is an enveloped, positive-sense and single-stranded RNA virus of about 30 kb. The life cycle of the virus with the host comprises mainly five steps as follows: attachment, penetration, biosynthesis, maturation and release. Once SARS-CoV-2 attaches to the host receptors, it penetrates the cells via endocytosis/membrane fusion. Herein, angiotensin-converting enzyme 2 is the entry and functional receptor of SARS-CoV-2. It has been shown that the spike for SARS-CoV-2, structural membrane proteins formed

by the *trans*-membrane trimetric glycoprotein protruding from the viral surface, also binds to angiotensin-converting enzyme 2. After the viral contents are released inside the host cells, viral RNA enters the nucleus to replicate. As for the biosynthesis, viral mRNA is used to make viral proteins. The new viral particles are formed in the maturation step and then released⁽²⁾.

Angiotensin-converting enzyme 2 plays an important role for the interaction between the classical and non-classical pathway of the renin angiotensin system. The former acts through the angiotensin II type 1 receptors, and its increased activity leads to fibrosis, inflammation and angiogenesis. The latter acts through the Mas receptors and has opposing effects to the angiotensin II type 1 receptors⁽³⁾. Angiotensin-converting enzyme 2 is expressed by the epithelial cells of lungs, intestines, kidneys and blood vessels; therefore, the aforementioned

Abbreviations: 25(OH)D, 25-hydroxyvitamin D; COVID-19, coronavirus disease; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

* **Corresponding author:** Timur Ekiz, email timurekiz@gmail.com




Fig. 1. The world map illustrates the total deaths and percentage of severe vitamin D deficiency in countries most commonly affected by COVID-19^(1,5-43). Severe vitamin D deficiency (%): (■), >30 (South Arabia, Qatar, Iran, China); (■), 20–30 (France, Chile, UK, Portugal); (■), 10–20 (Austria, Pakistan, Italy, Poland, Brazil, Israel, Croatia, Romania, Turkey, Germany); (■), 5–10 (India, Russia, Switzerland, Canada, Belgium, USA, South Korea, Ireland, Spain); (■), <5 (Greece, Singapore, Mexico, Japan, Ecuador, Australia, Sweden, Malaysia, Norway, Finland, Denmark, Netherlands). Total deaths: (●), >25 000 (USA, UK, Italy, France, Spain); (●), 5000–10 000 (Brazil, Belgium, Germany, Iran, The Netherlands, Canada); (●), 1000–5000 (China, Mexico, Turkey, Sweden, India, Ecuador, Russia, Peru, Switzerland, Ireland, Portugal, Romania); (●), 500–1000 (Poland, Pakistan, Japan, Austria, Denmark); (●), <500 (Chile, Finland, Saudi Arabia, Israel, South Korea, Norway, Greece, Malaysia, Australia, Croatia, Singapore, Qatar).

tissues/organs are vulnerable to SARS-CoV-2 infection⁽⁴⁾. Additionally, activation of the renin-angiotensin system is significantly associated with increased morbidity and mortality as in hypertension⁽³⁾.

On the other hand, vitamin D deficiency has also been described as pandemic and a global public health problem, especially in Europe (Table 1)⁽⁵⁻⁴³⁾. Regardless of age, ethnicity and latitude, recent data showed that 40% of Europeans are vitamin D deficient (25-hydroxyvitamin D (25(OH)D) levels <50 nmol/l), and 13% are severely deficient (25(OH)D <30 nmol/l)⁽⁴⁴⁾. According to regression analyses, a quadratic relationship was found between the prevalences of vitamin D deficiency in most commonly affected countries by COVID-19 and the latitudes (Fig. 2). Interestingly, vitamin D deficiency is more common in the subtropical and mid-latitude countries than the tropical and high-latitude countries. Contrary to the expectation, the most commonly affected countries with severe vitamin D deficiency are from the subtropical (Saudi Arabia 46%; Qatar 46%; Iran 33.4%; Chile 26.4%) and mid-latitude (France 27.3%; Portugal 21.2%; Austria 19.3%) regions. On the other hand, severe vitamin D deficiency was found to be nearly 0% in some high-latitude countries (e.g. Norway, Finland, Sweden, Denmark and Netherlands). The low prevalences of severe vitamin D deficiencies in high-latitude countries (except for the UK; 23.7%) can possibly be attributed to the high awareness of vitamin D deficiency, high amount of vitamin D supplementation, food fortification and health policies as well⁽⁴⁴⁾. Indeed, as the main source of vitamin D is exposure of the skin to sun (UV-B), it has long been supposed that living in a

sunny country guarantees sufficient vitamin D levels. However, there is increasing evidence that vitamin D deficiency may have been underestimated/ignored in low latitude, even in tropical countries⁽⁴⁵⁾.

The risks for vitamin D deficiency encompass obesity, elderly, lack of proper sun (UV-B) exposure, dark skin, smoking, living with air pollution and the presence of co-morbid diseases such as infection, cancer, CVD, chronic respiratory disease, osteoporosis, sarcopenia and diabetes mellitus^(46,47). Further, it is known that severe vitamin D deficiency dramatically increases the risk of mortality, infections and many other diseases. As such, it should indisputably be prevented whenever detected/possible⁽⁴⁶⁾.

Vitamin D hormone has important functions – including immunomodulatory, anti-inflammatory and anti-infective roles⁽⁴⁷⁾. It acts via monocyte and cell-mediated immunity stimulation, suppression of lymphocyte proliferation, antibody production and cytokine synthesis⁽⁴⁸⁾. Human lung cells are able to intracellularly convert the inactive 25(OH)D to its active form 1,25(OH)D which reduces proinflammatory cytokines and increases peptides (e.g. the innate antimicrobial peptide cathelicidin)⁽⁴⁸⁾. Cathelicidin has direct antiviral activity against enveloped respiratory viruses such as hepatitis B, influenza, respiratory syncytial virus and possibly the COVID-19 as well⁽⁴⁸⁾. Other than the above-mentioned functions, vitamin D has also anti-fibrotic effects. The renin-inhibiting activity and down-regulation of the renin-angiotensin system activity seem to be the beneficial effects of vitamin D. Moreover, vitamin D has been shown to suppress angiotensinogen and regulate its expression⁽⁴⁹⁾.

Table 1. Available data for vitamin D deficiency among older adults in countries most commonly affected by COVID-19

Rank*	Region/country	Latitude	n	Age (years): mean (±SD) or range	25(OH)D deficiency (%)		Reference and year
					<25 nmol/l†	<50 nmol/l	
Region of the Americas							
1	USA	Mid latitude (38°)	4363	76±5	7 (<30)	32	Eymundsdottir <i>et al.</i> ⁽⁵⁾ 2020
6	Brazil	Subtropics (23°)	908	73±5	14.4	58.0	Lopes <i>et al.</i> ⁽⁶⁾ 2014
13	Peru	Tropics (12°)	204	39±11	N/A	29.4	Pastor <i>et al.</i> ⁽⁸⁾ 2019
14	Canada	Mid latitude (45°)	11 336	3–79	7.4 (<30)	36.8	Sarafin <i>et al.</i> ⁽⁷⁾ 2015
18	Mexico	Tropics (19°)	585	41±15	2.0	43.6	Clark <i>et al.</i> ⁽¹⁰⁾ 2015
20	Chile	Subtropics (33°)	686	≥65	26.4 (<30)	64.9	Solis-Urra <i>et al.</i> ⁽¹¹⁾ 2019
21	Ecuador	Tropics (0°)	2374	71±8	N/A	21.6	Orces <i>et al.</i> ⁽⁹⁾ 2015
European region							
2	Spain	Mid latitude (41°)	312	≥85	14.4	N/A	Formiga <i>et al.</i> ⁽¹²⁾ 2014
3	Russia	High latitude (59°)	1664	3–75	8.0 (<30)	45.7	Karonova <i>et al.</i> ⁽¹⁷⁾ 2016
4	UK	High latitude (54°)	6004	<50	23.7 (<30)	55.3	Aspell <i>et al.</i> ⁽¹⁴⁾ 2019
5	Italy	Mid latitude (41°)	2640	65–98	10.6	21.6	Veronese <i>et al.</i> ⁽¹³⁾ 2014
7	France	Mid latitude (44°)	697	73±4	27.3	55.9	Cougnard-Grégoire <i>et al.</i> ⁽¹⁵⁾ 2015
8	Germany	High latitude (51°)	1671	60–85	10.7	48.6	Vetter <i>et al.</i> ⁽¹⁶⁾ 2020
9	Turkey	Mid latitude (37°)	1161	18–90	12.9	75.5	Öztürk <i>et al.</i> ⁽¹⁸⁾ 2017
15	Belgium	High latitude (50°)	697	32–53	7.3	51.1	Hoge <i>et al.</i> ⁽¹⁹⁾ 2015
17	Netherlands	High latitude (52°)	450	65–93	0.0	2.0	Ten Haff <i>et al.</i> ⁽²⁰⁾ 2019
22	Switzerland	Mid latitude (47°)	1291	≥60	8.0	39.2	Sakem <i>et al.</i> ⁽²¹⁾ 2013
23	Sweden	High latitude (56°)	995 (W)	80–81	0.0	16.0	Buchebner <i>et al.</i> ⁽²³⁾ 2014
24	Portugal	Mid latitude (38°)	3092	≥18	21.2	66.6	Duarte <i>et al.</i> ⁽²²⁾ 2020
27	Ireland	High latitude (53°)	1118	18–84	6.0	45.0	Cashman <i>et al.</i> ⁽²⁴⁾ 2013
28	Poland	High latitude (51°)	5775	16–90	16.0	65.8	Pludowski <i>et al.</i> ⁽²⁶⁾ 2016
30	Romania	Mid latitude (46°)	14 052	37–62	13.2	52.0	Niculescu <i>et al.</i> ⁽²⁷⁾ 2017
32	Austria	Mid latitude (47°)	161	65–80	19.3	64.0	Elmadfa <i>et al.</i> ⁽²⁵⁾ 2017
34	Denmark	High latitude (56°)	3409	19–72	0.0	23.6	Cashman <i>et al.</i> ⁽²⁸⁾ 2015
35	Norway	High latitude (69°)	12 817	30–87	0.3	18.6	Cashman <i>et al.</i> ⁽²⁹⁾ 2016
38	Finland	High latitude (64°)	4102	29–77	0.2	6.6	Cashman <i>et al.</i> ⁽²⁸⁾ 2015
39	Greece	Mid latitude (37°)	181 (M)	20–50	4.4	50.3	Kassi <i>et al.</i> ⁽³⁰⁾ 2015
40	Croatia	Mid latitude (45°)	120 (W)	61±9	14.2 (<30)	63.3	Laktasic <i>et al.</i> ⁽³¹⁾ 2010
Western Pacific Region							
11	China	Subtropics (31°)	2180	>65	30.6 (<30)	70.3	Wei <i>et al.</i> ⁽³²⁾ 2019
26	Singapore	Tropics (1°)	504	45–74	N/A	14.0	Robien <i>et al.</i> ⁽³³⁾ 2013
31	Japan	Mid latitude (38°)	9084	40–74	N/A	53.6	Nakamura <i>et al.</i> ⁽³⁴⁾ 2015
33	South Korea	Mid latitude (36°)	4107 (W)	50–79	6.6 (<37.5)	67.4	Shin <i>et al.</i> ⁽³⁵⁾ 2015
36	Australia	Subtropics (34°)	2413	51±17	0.9	22.7	Gill <i>et al.</i> ⁽³⁶⁾ 2014
37	Malaysia	Tropics (3°)	63 (M)	≥60	0.0	17.5	Chin <i>et al.</i> ⁽³⁷⁾ 2014
Eastern Mediterranean region							
10	Iran	Subtropics (30°)	370	≥35	33.4	61.9	Khosravi-Boroujeni <i>et al.</i> ⁽³⁸⁾ 2017
16	Saudi Arabia	Subtropics (24°)	3475	47±16	46.0	76.8	Alfawaz <i>et al.</i> ⁽³⁹⁾ 2014
19	Pakistan	Subtropics (30°)	858	18–60	N/A	58.4	Mehboobali <i>et al.</i> ⁽⁴⁰⁾ 2015
25	Qatar	Subtropics (25°)	547	49±13	46.0	N/A	El-Menyar <i>et al.</i> ⁽⁴²⁾ 2012
29	Israel	Subtropics (31°)	198 834	60 (median)	14.4	49.9	Saliba <i>et al.</i> ⁽⁴¹⁾ 2012
South-East Asia region							
12	India	Tropics (12°)	149	≥46	8.7 (<30)	48.3	Mechenro <i>et al.</i> ⁽⁴³⁾ 2018

n, Number; 25(OH)D, 25-hydroxyvitamin D; N/A, not applicable; W, women; M, men.
 * The most commonly infected countries and regions with COVID-19 in descending order.
 † Percentages of severe vitamin D deficiency.

The distribution of community outbreaks shows seasonal patterns along certain latitude, temperature and humidity, that is, similar to the behaviour of seasonal viral respiratory tract infections. It has been reported that COVID-19 displays significant spread in mid-latitude (35–50° N') regions and/or in those with an average temperature of 5–11°C and low humidity (Fig. 1)^(1,50). Coronaviruses are very stable at 4°C (viable for up to 3 d) and can survive at –20°C (for up to 2 years)⁽¹⁾. Depending on some parameters (e.g. temperature, humidity and sunlight), they can live on different surfaces for a few days. They are thermolabile; decreased sunlight, low temperatures and less humidity seem to be favourable for COVID-19⁽¹⁾. Although natural UV (UV-C) from the sunlight may not be strong

enough to kill COVID-19, its antimicrobial efficacy has long been shown to inactivate, thus preventing the transmission of airborne-mediated infections such as influenza and tuberculosis⁽⁵¹⁾. Further, UV-B from the sun can induce endogenous synthesis of vitamin D in the skin – being the main source of vitamin D other than the dietary intake or supplementation. These factors might possibly be explanatory as regards the low prevalence of COVID-19 in subtropical and southern countries.

Patients infected with COVID-19 have higher mortality rates if they are older, that is, 8.0% (70–79 years) and 14.8% (>80 years). The similar rates for co-morbid conditions are as 10.5% (CVD), 7.3% (diabetes mellitus), 6.3% (chronic




Fig. 2. The histogram shows the prevalence of vitamin D deficiency (<50 nmol/l) and severe deficiency (<25 nmol/l) among the forty countries most commonly affected by COVID-19. The number above each column represents the country's position in the world ranking concerning the number of total cases of infections. The colour band is a graphical representation of the four main climatic areas in the world. Regression lines show the prevalence of overall (solid black line) and severe (dotted line) vitamin D deficiencies. (■), Vitamin D deficiency; (■), severe vitamin D deficiency.

respiratory disease), 6.0% (hypertension) and 5.6% (cancer)⁽⁵²⁾. Older adults with any of these co-morbid diseases are at high risk for COVID-19 infection – especially in the presence of severe vitamin D deficiency⁽⁵³⁾. To this end, since there is positive/strong evidence concerning the effects of vitamin D against viral respiratory infections, it would not be unsound to say that vitamin D supplementation may decrease viral induction and inflammatory genes, and incidence/severity of respiratory tract infections⁽⁴⁸⁾. In this sense, a meta-analysis of twenty-five randomised controlled trials showed that vitamin D supplementation has a preventive effect against acute respiratory tract infections and that the benefit is higher in those subjects receiving daily or weekly vitamin D without additional bolus doses, and in those having severe vitamin D deficiency at baseline⁽⁵⁴⁾.

Although vitamin D was primarily recognised for bone metabolism, increasing evidence indicates its proper function for nearly every tissue in the body including brain, heart, lung, muscle, immune system and skin⁽⁵⁵⁾. Therefore, the treatment of vitamin D deficiency would be vital for several diseases including cardiovascular and neurological disorders, cancers, autoimmune diseases and infections as well⁽⁵⁵⁾. Likewise, a recent review recommended that in people at risk of influenza/COVID-19 infection, 250 µg/d of vitamin D₃ for a few weeks (or a month), that is, to rapidly increase the 25(OH)D concentrations and then 125 µg/d in the follow-up can be considered⁽⁴⁷⁾. The target should be to raise its value above 40–60 ng/ml. Additionally, the authors also suggested higher vitamin D₃ doses for infected patients with COVID-19. For sure, attention should be paid not to take high calcium supplementation for potential risk of hypercalcaemia while taking high doses of vitamin D₃. Needless to say, as vitamin D is synthesised mainly in the skin, sun (UV-B) exposure (15–20 min daily) inducing the light pink colour of minimal erythema would be the natural way of production and activation of vitamin D by keratinocytes⁽⁵⁵⁾.

Accordingly, presenting this paper, we would like to call attention to the possible association between severe vitamin D deficiency and mortality pertaining to COVID-19. Given its rare side effects and relatively wide safety, prophylactic vitamin D supplementation and/or food fortification might reasonably serve as a very convenient and incomparable/invaluable adjuvant therapy for these two worldwide public health problems alike.

Acknowledgements

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

M. K. performed study concept and design, acquisition of the data, writing and drafting the manuscript, and final approval and is the guarantor of the manuscript. T. E. performed study concept and design, acquisition of the data, writing and drafting the manuscript and final approval. V. R. performed study concept and design, writing and drafting the manuscript and final approval. Ö. K. performed study concept and design, writing and drafting the manuscript and final approval. K.-V. C. performed study concept and design, writing and drafting the manuscript and final approval. L. Ö. performed study concept and design, writing and drafting the manuscript, and final approval and is the supervisor.

The authors report no conflicts of interest.

References

1. World Health Organization (2020) Coronavirus disease (COVID-2019) situation reports. Geneva: World Health Organization. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/> (accessed May 2020).

2. Yuki K, Fujiogi M & Koutsogiannaki S (2020) COVID-19 pathophysiology: a review. *Clin Immunol* **2020**, 108427.
3. Vaduganathan M, Vardeny O, Michel T, *et al.* (2020) Renin-angiotensin-aldosterone system inhibitors in patients with Covid-19. *N Engl J Med* **382**, 1653–1659.
4. Zou X, Chen K, Zou J, *et al.* (2020) Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* (epublication ahead of print version 12 March 2020).
5. Eymundsdottir H, Chang M, Geirsdottir OG, *et al.* (2020) Lifestyle and 25-hydroxy-vitamin D among community-dwelling old adults with dementia, mild cognitive impairment, or normal cognitive function. *Aging Clin Exp Res* (epublication ahead of print version 4 April 2020).
6. Lopes JB, Fernandes GH, Takayama L, *et al.* (2014) A predictive model of vitamin D insufficiency in older community people: from the Sao Paulo Aging & Health study (SPAHS). *Maturitas* **78**, 335–340.
7. Sarafin K, Durazo-Arvizu R, Tian L, *et al.* (2015) Standardizing 25-hydroxyvitamin D values from the Canadian Health Measures Survey. *Am J Clin Nutr* **102**, 1044–1050.
8. Urrunaga-Pastor D, Guarnizo-Poma M, Macollunco-Flores P, *et al.* (2019) Association between vitamin D deficiency and insulin resistance markers in euthyroid non-diabetic individuals. *Diabetes Metab Syndr* **13**, 258–263.
9. Orces CH (2015) Vitamin D status among older adults residing in the littoral and Andes mountains in Ecuador. *ScientificWorldJournal* **2015**, 545297.
10. Clark P, Vivanco-Muñoz N, Piña JT, *et al.* (2015) High prevalence of hypovitaminosis D in Mexicans aged 14 years and older and its correlation with parathyroid hormone. *Arch Osteoporos* **10**, 225.
11. Solis-Urra P, Cristi-Montero C, Romero-Parra J, *et al.* (2019) Passive commuting and higher sedentary time is associated with vitamin D deficiency in adult and older women: results from Chilean National Health Survey 2016–2017. *Nutrients* **11**, 300.
12. Formiga F, Ferrer A, Megido MJ, *et al.* (2014) Low serum vitamin D is not associated with an increase in mortality in oldest old subjects: the Octabaix three-year follow-up study. *Gerontology* **60**, 10–15.
13. Veronese N, Sergi G, De Rui M, *et al.* (2014) Serum 25-hydroxyvitamin D and incidence of diabetes in elderly people: the PRO.V.A. Study. *J Clin Endocrinol Metab* **99**, 2351–2358.
14. Aspell N, Laird E, Healy M, *et al.* (2019) The prevalence and determinants of vitamin D status in community-dwelling older adults: results from the English Longitudinal Study of Ageing (ELSA). *Nutrients* **11**, 1253.
15. Cougnard-Grégoire A, Merle BM, Korobelnik JF, *et al.* (2015) Vitamin D deficiency in community-dwelling elderly is not associated with age-related macular degeneration. *J Nutr* **145**, 1865–1872.
16. Vetter VM, Spira D, Banszerus VL, *et al.* (2020) Epigenetic clock and leukocyte telomere length are associated with vitamin D status, but not with functional assessments and frailty in the Berlin Aging Study II. *J Gerontol A Biol Sci Med Sci* (epublication ahead of print version 23 April 2020).
17. Karonova T, Andreeva A, Nikitina I, *et al.* (2016) Prevalence of vitamin D deficiency in the North-West region of Russia: a cross-sectional study. *J Steroid Biochem Mol Biol* **164**, 230–234.
18. Öztürk ZA, Gol M & Türkbeyler İH (2017) Prevalence of vitamin D deficiency in otherwise healthy individuals between the ages of 18 and 90 years in southeast Turkey. *Wien Klin Wochenschr* **129**, 854–855.
19. Hoge A, Donneau AF, Strel S, *et al.* (2015) Vitamin D deficiency is common among adults in Wallonia (Belgium, 51°30' North): findings from the Nutrition, Environment and Cardio-Vascular Health study. *Nutr Res* **35**, 716–725.
20. Ten Haaf DSM, Balvers MGJ, Timmers S, *et al.* (2019) Determinants of vitamin D status in physically active elderly in the Netherlands. *Eur J Nutr* **58**, 3121–3128.
21. Sakem B, Nock C, Stanga Z, *et al.* (2013) Serum concentrations of 25-hydroxyvitamin D and immunoglobulins in an older Swiss cohort: results of the Senior Labor Study. *BMC Med* **11**, 176.
22. Duarte C, Carvalheiro H, Rodrigues AM, *et al.* (2020) Prevalence of vitamin D deficiency and its predictors in the Portuguese population: a nationwide population-based study. *Arch Osteoporos* **15**, 36.
23. Buchebner D, McGuigan F, Gerdhem P, *et al.* (2014) Vitamin D insufficiency over 5 years is associated with increased fracture risk-an observational cohort study of elderly women. *Osteoporos Int* **25**, 2767–2775.
24. Cashman KD, Kiely M, Kinsella M, *et al.* (2013) Evaluation of vitamin D standardization program protocols for standardizing serum 25-hydroxyvitamin D data: a case study of the program's potential for national nutrition and health surveys. *Am J Clin Nutr* **97**, 1235–1242.
25. Elmadfa I, Meyer AL, Wottawa D, *et al.* (2017) Vitamin D intake and status in Austria and its effects on some health indicators. *Austin J Nutr Metab* **4**, 1050.
26. Phudowski P, Ducki C, Konstantynowicz J, *et al.* (2016) Vitamin D status in Poland. *Pol Arch Med Weun* **126**, 530–539.
27. Niculescu DA, Capatina CAM, Dusceac R, *et al.* (2017) Seasonal variation of serum vitamin D levels in Romania. *Arch Osteoporos* **12**, 113.
28. Cashman KD, Dowling KG, Škrabáková Z, *et al.* (2015) Standardizing serum 25-hydroxyvitamin D data from four Nordic population samples using the vitamin D standardization program protocols: shedding new light on vitamin D status in Nordic individuals. *Scand J Clin Lab Invest* **75**, 549–561.
29. Cashman KD, Dowling KG, Škrabáková Z, *et al.* (2016) Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr* **103**, 1033–1044.
30. Kassi EN, Stavropoulos S, Kokkoris P, *et al.* (2015) Smoking is a significant determinant of low serum vitamin D in young and middle-aged healthy males. *Hormones (Athens)* **14**, 245–250.
31. Laktasic-Zerjavic N, Korsic M, Crncevic-Orlic Z, *et al.* (2010) Vitamin D status, dependence on age, and seasonal variations in the concentration of vitamin D in Croatian postmenopausal women initially screened for osteoporosis. *Clin Rheumatol* **29**, 861–867.
32. Wei J, Zhu A & Ji JS (2019) A comparison study of vitamin D deficiency among older adults in China and the United States. *Sci Rep* **9**, 19713.
33. Robien K, Butler LM, Wang R, *et al.* (2013) Genetic and environmental predictors of serum 25-hydroxyvitamin D concentrations among middle-aged and elderly Chinese in Singapore. *Br J Nutr* **109**, 493–502.
34. Nakamura K, Kitamura K, Takachi R, *et al.* (2015) Impact of demographic, environmental, and lifestyle factors on vitamin D sufficiency in 9084 Japanese adults. *Bone* **74**, 10–17.
35. Shin JH, Lee HT, Lim YH, *et al.* (2015) Defining vitamin D deficiency and its relationship to hypertension in postmenopausal Korean women. *J Women's Health (Larchmt)* **24**, 1021–1029.

36. Gill TK, Hill CL, Shanahan EM, *et al.* (2014) Vitamin D levels in an Australian population. *BMC Public Health* **14**, 1001.
37. Chin KY, Ima-Nirwana S, Ibrahim S, *et al.* (2014) Vitamin D status in Malaysian men and its associated factors. *Nutrients* **6**, 5419–5433.
38. Khosravi-Boroujeni H, Sarrafzadegan N, Sadeghi M, *et al.* (2017) Prevalence and trends of vitamin D deficiency among Iranian adults: a longitudinal study from 2001–2013. *J Nutr Sci Vitaminol (Tokyo)* **63**, 284–290.
39. Alfawaz H, Tamim H, Alharbi S, *et al.* (2014) Vitamin D status among patients visiting a tertiary care center in Riyadh, Saudi Arabia: a retrospective review of 3475 cases. *BMC Public Health* **14**, 159.
40. Mehboobali N, Iqbal SP & Iqbal MP (2015) High prevalence of vitamin D deficiency and insufficiency in a low income peri-urban community in Karachi. *J Pak Med Assoc* **65**, 946–949.
41. Saliba W, Rennert HS, Kershenbaum A, *et al.* (2012) Serum 25(OH)D concentrations in sunny Israel. *Osteoporos Int* **23**, 687–694.
42. El-Menyar A, Rahil A, Dousa K, *et al.* (2012) Low vitamin D and cardiovascular risk factors in males and females from a sunny, rich country. *Open Cardiovasc Med J* **6**, 76–80.
43. Mechenro J, Venugopal G, Kumar B, *et al.* (2018) Vitamin D status in Kancheepuram District, Tamil Nadu, India. *BMC Public Health* **18**, 1345.
44. Lips P, Cashman KD, Lamberg-Allardt C, *et al.* (2019) Current vitamin D status in European and Middle East countries and strategies to prevent vitamin D deficiency: a position statement of the European Calcified Tissue Society. *Eur J Endocrinol* **180**, P23–P54.
45. Mendes MM, Hart KH, Botelho PB, *et al.* (2018) Vitamin D status in the tropics: is sunlight exposure the main determinant? *Nutr Bull* **43**, 428–434.
46. Schleicher RL, Sternberg MR, Looker AC, *et al.* (2016) National estimates of serum total 25-hydroxyvitamin D and metabolite concentrations measured by liquid chromatography-tandem mass spectrometry in the US population during 2007–2010. *J Nutr* **146**, 1051–1061.
47. Grant WB, Lahore H, McDonnell SL, *et al.* (2020) Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients* **12**, E988.
48. Beard JA, Bearden A & Striker R (2011) Vitamin D and the antiviral state. *J Clin Virol* **50**, 194–200.
49. Feldman D, Pike JW, Bouillon R, *et al.* (editors) (2017) *Vitamin D: Volume 1: Biochemistry, Physiology And Diagnostics*. Cambridge: Academic Press.
50. Sajadi MM, Habibzadeh P, Vintzileos A, *et al.* (2020) Temperature, humidity and latitude analysis to predict potential spread and seasonality for COVID-19. <https://ssrn.com/abstract=3550308> (accessed March 2020).
51. Welch D, Buonanno M, Grilj V, *et al.* (2018) Far-UVC light: a new tool to control the spread of airborne-mediated microbial diseases. *Sci Rep* **8**, 2752.
52. Novel Coronavirus Pneumonia Emergency Response Epidemiology Team (2020) The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi* **41**, 145–151.
53. Alipio M (2020) Vitamin D supplementation could possibly improve clinical outcomes of patients infected with Coronavirus-2019 (COVID-2019). <https://ssrn.com/abstract=3571484> (accessed April 2020).
54. Martineau AR, Jolliffe DA, Hooper RL, *et al.* (2017) Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* **356**, i6583.
55. Mostafa WZ & Hegazy RA (2015) Vitamin D and the skin: focus on a complex relationship: a review. *J Adv Res* **6**, 793–804.